

THE OPTIMIST CLASSES

IIT-JAM TOPPERS

8
AIR



MANOJ KUMAR SINGH

9
AIR



ABHAY

12
AIR



PAWAN

15
AIR



SATYAM

21
AIR



SOUMIL GIRISH SAHU

29
AIR




BHOOMIJA

31
AIR



AKSHIT AGGARWAL

32
AIR



SHIKHAR CHAMOLI

33
AIR



RAVI SINGH ADHIKARI

44
AIR



GAURAV JHA

56
AIR



SWAPNIL JOSHI

65
AIR



LOKESH BHATT

75
AIR



GOPESH VISHVAKARMA

77
AIR



VAIBHAV

80
AIR



SHASWAT CHAMOLI

CSIR-NET-JRF RESULTS 2022



ANNU
DL01000308



ALANKAR
UP15000162



SAHIL RANA
HR09000108



JAYESTHI
RJ11000161



DASRATH
RJ06000682



VIVEK
UK01000439



UZAIR AHMED
UP02000246



SURYA PRATAP SINGH
RJ06000232



HIMANSHU
UP10000095



CHANDAN
RJ09000159



SAIKHOM JOHNSON
MN01000196



AJAY SAINI
RJ06001744



VIKAS YADAV
RJ06001102



JYOTSNA KOHLI
UK02000262



SHYAM SUNDAR
RJ06000615

THE OPTIMIST CLASSES

AN INSTITUTE FOR NET-JRF/GATE/IIT-JAM/JEST/TIFR/M.Sc ENTRANCE EXAMS

CONTACT: 9871044043

GATE PAPER 2003


Q. 1- Q.30: Carry ONE mark each

- Q1. The two vectors $P = \hat{i}$, $q = (i + \hat{j})/\sqrt{2}$ are
 (a) related by a rotation (b) related by a reflection through xy - plane
 (c) related by an inversion (d) not linearly independent
- Q2. A 3×3 matrix has eigenvalues $0, 2 + i, 2 - i$. Which of the following statements is correct?
 (a) The matrix is Hermitian (b) The matrix is unitary
 (c) The inverse of the matrix exists (d) The determinant of the matrix is zero.
- Q3. The value of the integral $\int_C dz / z^2$, where z is a complex variable and C is the unit circle with the origin as its centre, is:
 (a) 0 (b) $2\pi i$ (c) $4\pi i$ (d) $-4\pi i$
- Q4. A particle with an initial velocity $v_0 \hat{i}$ enters a region with an electric field $E_0 \hat{j}$ and a magnetic field $B_0 \hat{j}$. The trajectory of the particle will
 (a) be an ellipse (b) be a cycloid
 (c) be a helix with constant pitch (d) not be confined to any plane
- Q5. An object of mass m rests on a surface with coefficient of static friction μ . Which of the following is NOT correct?
 (a) The force of friction is exactly μmg
 (b) The maximum force of friction is μmg
 (c) The force of friction is along the surface
 (d) The force of friction opposes any effort to move the object
- Q6. The Lagrangian of a particle of mass m moving in a plane is given by $L = (1/2)m[v_x^2 + v_y^2] + a(xv_y - yv_x)$, where v_x and v_y are velocity components and a is a constant. The canonical momenta of the particle are given by
 (a) $p_x = mv_x$ and $p_y = mv_y$ (b) $p_x = mv_x + ay$ and $p_y = mv_y + ax$
 (c) $p_x = mv_x - ay$ and $p_y = mv_y + ax$ (d) $p_x = mv_x - ay$ and $p_y = mv_y - ax$
- Q7. Two events are separated by a distance of 6×10^5 km and the first event occurs 1 sec before the second event. The interval between the two events
 (a) is time-like (b) is light-like (null)
 (c) is space-like (d) cannot be determined from the information given

- Q8. An electric charge, $+Q$ is placed on the surface of a solid, conduction sphere of radius a . The distance measured from the centre of the sphere is denoted as r . Then:
- The charge gets distributed uniformly through the volume of the sphere
 - The electrostatic potential has the same value for $r < a$
 - An equal and opposite charge gets induced in the bottom half of the sphere
 - The electric field is given by $1/(4\pi\epsilon_0 r^2)$ for $r < a$
- Q9. An electric field applied along the length of a long cylinder produces a polarization P . The depolarization field produced in this configuration is :
- $4\pi P/3$
 - $-4\pi P/3$
 - $2\pi P$
 - 0
- Q10. Which one of the following Maxwell's equations implies the absence of magnetic monopoles?
- $\vec{\nabla} \cdot \vec{E} = \rho / \epsilon_0$
 - $\vec{\nabla} \cdot \vec{B} = 0$
 - $\vec{\nabla} \times \vec{E} = -\partial B / \partial t$
 - $\vec{\nabla} \times \vec{B} = (1/c^2)\partial \vec{B} / \partial t + \mu_0 \vec{J}$
- Q11. An electromagnetic wave is propagating in free space in the z -direction. If the electric field is given by $E = \cos(\omega t - kz) \hat{i}$, where $\omega t = ck$, then the magnetic field is given by :
- $B = (1/c)\cos(\omega t - kz) \hat{j}$
 - $B = (1/c)\sin(\omega t - kz) \hat{j}$
 - $B = (1/c)\cos(\omega t - kz) \hat{i}$
 - $B = (1/c)\sin(\omega t - kz) \hat{i}$
- Q12. Given a wave with the dispersion relation $\omega = ck + m$ for $k > 0$ and $m > 0$, which one of the following is true?
- The group velocity is greater than the phase velocity
 - The group velocity is less than the phase velocity
 - The group velocity and the phase velocity are equal
 - There is no definite relation between the group velocity and the phase velocity
- Q13. Which of the following is a valid normalized wave function for a particle in a one dimensional infinite potential well of width L centred at $x = 0$?
- $(2/L)[\cos(2\pi x/L) + \sin(2\pi x/L)]$
 - $(2/L)^{1/2} \sin[n\pi x/L]$ for odd n
 - $(2/L)^{1/2} \cos[n\pi x/L]$ for odd n
 - $(2/L)\cos(\pi x/L)$
- Q14. The commutator $[x, P^2]$, where x and P are position and momentum operators respectively, is
- $2i\hbar P$
 - $-i\hbar P$
 - $2i\hbar xP$
 - $-2i\hbar xP$
- Q15. A spin half particle is in the state $S_z = \frac{\hbar}{2}$. The expectation values of S_x, S_x^2, S_y, S_y^2 are given by
- $0, 0, \hbar^2/4, \hbar^2/4$
 - $0, \hbar^2/4, \hbar^2/4, 0$
 - $0, \hbar^2/4, 0, \hbar^2/4$
 - $\hbar^2/4, \hbar^2/4, 0, 0$
- Q16. The spectral term for the atom with 70% subshell and only $S = 3/2$ is
- 3P_0
 - ${}^4F_{9/2}$
 - ${}^3F_{1/2}$
 - ${}^4P_{1/2}$
- Q17. The hyperfine splitting of the spectral lines of an atom is due to
- the coupling between the spins of two or more electrons
 - the coupling between the spins and the orbital angular momenta of the electrons
 - the coupling between the electron spins and the nuclear spin
 - the effect of external electromagnetic fields
- Q18. A piston containing an ideal gas is originally in the state X (see figure). The gas is taken through a thermal cycle $X \rightarrow Y \rightarrow X$ as shown:

The work done by the gas is positive if the direction of the thermal cycle is :

- (a) clockwise
- (b) counter - clockwise
- (c) neither clockwise nor counter - clockwise
- (d) clockwise from $X \rightarrow Y$ and counter - clockwise from $Y \rightarrow X$.



Q19. A second order phase transition is one in which

- (a) the plot of entropy as a function of temperature shows a discontinuity.
- (b) the plot of specific heat as a function of temperature shows a discontinuity.
- (c) the plot of volume as a function of pressure shows a discontinuity.
- (d) the plot of compressibility as a function of temperature is continuous.

Q20. Consider the Fermi-Dirac distribution function $f(E)$ at room temperature ($300K$) where E refers to energy. If

E_F is the Fermi energy, which of the following is true?

- (a) $f(E)$ is a step function
- (b) $f(E_F)$ has a value of $1/2$
- (c) States with $E < E_F$ are filled completely
- (d) $f(E)$ is large and tends to infinity as E decreases much below E_F .

Q21. If the ionic radii of Mn and S are 0.80 and $0.184nm$ respectively, the structure of MnS will be

- (a) cubic closed packed
- (b) body centered cubic
- (c) NaCl type
- (d) primitive cubic cell

Q22. A cubic cell consists of two atoms of masses m_1 and m_2 ($m_1 > m_2$) with m_1 and m_2 atoms situated on alternate planes. Assuming only nearest neighbor interactions, the centre of mass of the two atoms

- (a) moves with the atoms in the optical mode and remains fixed in the acoustic mode
- (b) remains fixed in the optical mode and moves with the atoms in the acoustic mode
- (c) remains fixed in both optical and acoustic modes
- (d) moves with the atoms in both optical and acoustic modes

Q23. In simple metals the phonon contribution to the electrical resistivity at temperature T is

- (a) directly proportional to T above Debye temperature and to ' T^3 ' below it
- (b) inversely proportional to T for all temperatures
- (c) independent of T for all temperature
- (d) directly proportional to T above Debye temperature and to T^5 below it

Q24. The effective mass of an electron in a semiconductor can be

- (a) negative near the bottom of the band
- (b) a scalar quantity with a small magnitude
- (c) zero at the center of the band
- (d) negative near the top of the band

Q25. The dielectric constant of water is 80 . However its refractive index is 1.75 invalidating the expression $n = \epsilon^{1/2}$. This is because

- (a) the water molecule has a permanent dipole moment
- (b) the boiling point of water is $100^\circ C$
- (c) the two quantities are measured in different experiments
- (d) water is transparent to visible light

Q26. The nucleus of the atom 9Be_4 consists of

- (a) 13 up quarks and 13 down quarks
- (b) 13 up quarks and 14 down quarks
- (c) 14 up quarks and 13 down quarks
- (d) 14 up quarks and 14 down quarks

Q27. Which one of the following nuclear reactions is possible?

- (a) ${}^{14}N_7 \rightarrow {}^{13}C_6 + \beta^+ + \nu_e$
- (b) ${}^{13}N_7 \rightarrow {}^{13}C_6 + \beta^+ + \nu_e$

(c) ${}^{13}\text{N}_7 \rightarrow {}^{13}\text{C}_6 + \beta^+$

(d) ${}^{13}\text{N}_7 \rightarrow {}^{13}\text{C}_7 + \beta^+ + \nu_e$

- Q28. Suppose that a neutron at rest in free space decays into a proton and an electron. This process would violate
- (a) conservation of charge (b) conservation of energy
(c) conservation of linear momentum (d) conservation of angular momentum

- Q29. Which one of the following is TRUE for a semiconductor pn junction with no external bias?
- (a) The total charge in the junction is not conserved
(b) The p side of the junction is positively charged
(c) The p side of the junction is negative charged
(d) No charge develops anywhere in the junction

- Q30. Which one of the set of values given below does NOT satisfy the Boolean relation $R = PQ'$ (where Q' denotes NOT Q)?

(a) $P=1, Q=1, R=0$ (b) $P=1, Q=1, R=1$ (c) $P=0, Q=0, R=0$ (d) $P=0, Q=1, R=1$

Q.31 - Q.90: Carry ONE mark each.

- Q31. The curl of the vector $A = zi + xj + yk$ is given by

(a) $i + j + k$ (b) $i - j + k$ (c) $i + j - k$ (d) $-i - j - k$

- Q32. Consider the differential equation $d^2x/dt^2 + 2dx/dt + x = 0$. At time $t = 0$, it is given that $x = 1$ and $dx/dt = 0$. At $t = 1$, the value of x is given by:

(a) $1/e$ (b) $2/e$ (c) 1 (d) $3/e$

- Q33. S_{ij} and A_{ij} represent a symmetric and an antisymmetric real-valued tensor respectively in three dimensions. The number of independent components of S_{ij} and A_{ij} are

(a) 3 and 6 respectively (b) 6 and 3 respectively
(c) 6 and 6 respectively (d) 9 and 6 respectively

- Q34. Consider the four statements given below about the function $f(x) = x^4 - x^2$ in the range $-\infty < x < +\infty$.

Which one of the following statements is correct?

P the plot of $f(x)$ versus x has two maxima and two minima

Q the plot of $f(x)$ versus x cuts the x -axis at four points

R the plot of $f(x)$ versus x has three extrema

S no part of the plot $f(x)$ versus x lies in the fourth quadrant

Pick the right combination of correct choices from those given below

(a) P and R (b) R only (c) R and S (d) P and Q

- Q35. The Fourier transform of the function $f(x)$ is $F(k) = \int e^{ikx} f(x) dx$. The Fourier transform of $df(x)/dx$ is

(a) $dF(k)/dk$ (b) $\int F(k)/dk$ (c) $-ikF(k)$ (d) $ikF(k)$

- Q36. A particle of mass m is moving in a potential of the form $V(x, y, z) = (1/2)m\omega^2(3x^2 + 3y^2 + 2z^2 + 2xy)$.

The oscillation frequencies of the three normal modes of the particle are given by

(a) $\omega, \sqrt{3}\omega$ and $\sqrt{3}\omega$ (b) $\sqrt{2}\omega, \sqrt{3}\omega$ and $\sqrt{3}\omega$

(c) $\sqrt{2}\omega, \sqrt{2}\omega$ and 2ω (d) $\sqrt{2}\omega, 2\omega$ and 2ω

Q37. The speed of a particle whose kinetic energy is equal to its rest mass energy is given by (c is the speed of light in vacuum)

- (a) $c/3$ (b) $\sqrt{2}c/3$ (c) $c/2$ (d) $\frac{\sqrt{3}}{2}c$

Q38. Electromagnetic waves are propagating along a hollow, metallic waveguide whose cross-section is a square of side W . The minimum frequency of the electromagnetic waves is

- (a) c/W (b) $2c/W$ (c) $\pi c/W$ (d) $\sqrt{2}\pi c/W$

Q39. Consider the given statements about $E(r, t)$ and $B(r, t)$, The electric and magnetic vectors respectively in a region of free space

- P** Both E and B are conservative vector fields
Q Both E and B are central force fields
R E and B are mutually perpendicular in the region
S Work done by B on a moving charge in the region is zero

Choose the right combination of correct statements from the following :

- (a) P and R (b) R and S (c) S only (d) P and Q

Q40. An infinite conducting sheet in the x - y plane carries a surface current density K along the y -axis. The magnetic field B for $z > 0$ is :

- (a) $B = 0$ (b) $B = \mu_0 K \hat{k} / z$ (c) $B = \mu_0 K \hat{i} / 2$ (d) $B = \mu_0 K \hat{j} / (x^2 + z^2)^{0.5}$

Q41. A parallel beam of infrared radiation of wavelength of $1.01 \times 10^{-6} m$ is incident normally on a screen with two slits $5 \times 10^{-6} m$ apart and the resulting interference pattern is observed on a distant screen. What is the largest number of maxima that can be observed on the screen?

- (a) 4 (b) 9 (c) 13 (d) infinitely many

Q42. A parallel beam of electrons of a given momentum pass through a screen S_1 containing a slit and produces a diffraction pattern on a screen S_2 , placed behind it. The width of the central maximum observed on the screen S_2 can be increased by

- (a) decreasing the distance between the screen S_1 and S_2
 (b) increasing the width of the slit in screen S_1 .
 (c) decreasing the momentum of the electrons.
 (d) increasing the momentum of the electrons.

Q43. An electron in a time independent potential is in a state which is a superposition of the ground state ($E_0 = 11 eV$) and the first excited state ($E_1 = 13 eV$). The wave function of the electron will repeat itself with a period of

- (a) 3.1×10^{-18} sec (b) 2.1×10^{-15} sec (c) 1.2×10^{-12} sec (d) 1.0×10^{-9} sec

Q44. A particle has the wave function $\psi(x, t) = A \exp(i\omega t) \cos(kx)$. Which one of the following is correct?

- (a) This is an eigen state of both energy and momentum.
 (b) This is an eigen state of momentum and not energy.
 (c) This is an eigen state of energy and not momentum.
 (d) This is not an eigen state of energy or momentum.

Q45. A free particle with energy E whose wave-function is a plane wave with wavelength λ enters a region of constant potential $V > 0$ where the wavelength of the particle is 2λ . The ratio (V/E) is

- (a) $1/2$ (b) $2/3$ (c) $3/4$ (d) $4/5$

Q46. The vibrational spectrum of a molecule exhibits a strong line with P and R branches at a frequency ν_1 and a

weaker line at a frequency ν_2 . The frequency ν_3 is not shown up. Its vibrational Raman spectrum shows a strongly polarized line at frequency ν_3 and no feature at ν_1 and ν_2 .

- (a) the molecule could be linear (b) the molecule lacks a center of inversion
(c) ν_1 arises from a symmetric stretching mode (d) ν_3 arises from a bending mode

Q47. Three value of rotational energies of molecules are given below in different units

P 10cm^{-1}

Q 10^{-23}J

R 10^4MHz

Choose the correct arrangement in the increasing order of energy

- (a) P, Q, R (b) R, Q, P (c) R, P, Q (d) Q, R, P

Q48. The short wavelength cut off of the continuous X-ray spectrum from a nickel target is 0.0825nm . The voltage required to the applied to an X-ray tube is

- (a) 0.15KV (b) 1.5KV (c) 15KV (d) 150KV

Q49. The spin-orbit coupling constant for the upper state of sodium atom which emits D lines of wave numbers 16956.2 and 16973.4cm^{-1} is:

- (a) 15cm^{-1} (b) 11.4cm^{-1} (c) 12.5cm^{-1} (d) 15.1cm^{-1}

Q50. Consider the following statements about molecular spectra

P CH_4 does not give pure rotational Raman lines

Q SF_6 could be studied by rotational Raman spectroscopy

R N_2 shows infrared absorption spectrum

S CH_3CH_3 shows vibrational Raman and infrared absorption lines

T H_2O_2 shows pure rotational spectrum

Choose the right combination of correct statements

- (a) P and Q (b) P, R and T (c) P, S and T (d) Q and R

Q51. The temperature of a cavity of fixed volume is double. Which of the following is true for the black-body radiation inside the cavity?

- (a) its energy and the number of photons both increase 8 times
(b) its energy increases 8 times and the number of photons 16 times
(c) its energy increases 16 times and the number of photons increases 8 times
(d) its energy and the number of photons both increase 16 times

Q52. A sample of ideal gas with initial pressure P and volume V is taken through an isothermal expansion proceed during which the change in entropy is found to be ΔS . The universal gas constant is R . Then the work done by the gas is given by

- (a) $(PV\Delta S)/(nR)$ (b) $nR\Delta S$ (c) PV (d) $(P\Delta S)/(nRV)$

Q53. Hydrogen molecules (mass m) are in thermal equilibrium at a temperature T . Assuming classical distribution of velocity, the most probable speed at room temperature is:

- (a) $(k_B T)/m$ (b) $2k_B T/m$ (c) $(\sqrt{2k_B T}/m)$ (d) $m/(\sqrt{2k_B T})$

Q54. Consider the energy E in the first Brillouin zone as a function of the magnitude of the wave vector k for a crystal of lattice constant a . Then

- (a) the slope of E versus k is proportional to the group velocity
(b) the slope of E versus k has its maximum value at $|k| = \pi/a$
(c) the plot of E versus k will be parabolic in the interval $(-\pi/a) < k < (\pi/a)$
(d) the slope of E versus k is non-zero for all k the interval $(-\pi/a) < k < (\pi/a)$

- Q55. An external magnetic field of magnitude H is applied to a Type-I superconductor at a temperature below the transition point. Then which one of the following statements is NOT true for H less than the critical field H_c ?
- the sample is diamagnetic
 - its magnetization varies linearly with H
 - the lines of magnetic induction are pushed out from the sample
 - the sample exhibits mixed states of magnetization near H_c
- Q56. A ferromagnetic material has a Curie temperature $100K$. Then
- its susceptibility is doubled when it is cooled from $300K$ to $200K$
 - all the atomic magnets in it get oriented in the same direction above $100K$
 - the plot of inverse susceptibility versus temperature is linear with a slope T_c
 - The plot of its susceptibility versus temperature is linear with an intercept T_c

- Q57. The point group symmetries of the three molecules shown in Figs, 1-3 are respectively




Fig.1




Fig.2




Fig. 3

[notation : $C_{2v} = 2mm$;

$C_{2h} = 2/m$

$D_{2h} = mmm$]

- C_{2h}, C_{2v}, C_{2h}
 - C_{2v}, C_{2h}, C_{2h}
 - D_{2h}, C_{2v}, C_{2h}
 - C_{2v}, D_{2h}, C_{2h}
- Q58. The energy density of states of an electron in a one dimensional potential well of infinitely high walls is (the symbols have their usual meaning)

(a) $L\sqrt{m} / \left[\pi\hbar\sqrt{(2E)} \right]$

(b) $Lm / \left(\pi\hbar\sqrt{E} \right)$

(c) $Lm / \left[\pi\hbar\sqrt{(2E)} \right]$


(d) $L\sqrt{m} / (2\pi\hbar E)$

- Q59. Which one of the following statements concerning the Compton effect is NOT correct?
- The wavelength of the scattered photon is greater than or equal to the wavelength of the incident photon.
 - The electron can acquire a kinetic energy equal to the energy of the incident photon.
 - The energy of the incident photon equals to the kinetic energy of the electron plus the energy of the scattered photon.
 - The kinetic energy acquired by the electron is the largest when the incident and scattered photons move in opposite directions.
- Q60. If the photon were to have a finite mass, then the Coulomb potential between two stationary charges separated by a distance r would
- be strictly zero beyond some distance
 - fall off exponentially for large values of r
 - fall off as $1/r^3$ for large values of r
 - fall off as $1/r$ for large values of r

- Q61. A stationary particle in free space is observed to spontaneously decay into two photons. This implies that
- the particle carries electric charge
 - the spin of the particle must be greater than or equal to 2
 - the particle is a boson
 - the mass of the particle must be greater than or equal to the mass of the hydrogen atom

- Q62. The masses of hydrogen atom, neutron and $^{238}\text{U}_{92}$ are given by 1.0078, 1.0087 and 238.0508 respectively. The binding energy of $^{238}\text{U}_{92}$ is therefore approximately equal to (Taking 1 a.m.u. \equiv 931.64 MeV)
- 120 MeV
 - 1500 MeV
 - 1600 MeV
 - 1800 MeV

- Q63. A bistable multivibrator with a saturation voltage $\pm 5V$ is shown in the diagram. The positive and negative threshold at the inverting terminal for which the multivibrator will switch to the other state are



- (a) $\pm 5/11V$ (b) $\pm 10/11V$ (c) $\pm 5V$ (d) $\pm 11V$
- Q64. An Avalanche effect is observed in a diode when
- (a) the forward voltage is less than the breakdown voltage
 (b) the forward voltage exceeds the breakdown voltage
 (c) the reverse voltage exceeds the breakdown voltage
 (d) the diode is heavily doped and forward biased

- Q65. Which of the given relations between the Boolean variables P and Q is NOT correct? (In the notation used here, P' denotes NOT P and Q' denotes NOT Q)

(a) $PQ' + PQ = P$ (b) $(PQ)' + P' + Q'$ (c) $PQ' + (P' + Q)'$ (d) $PQ' + Q = P$

Data for Q. No. 66 and Q.67


Consider the vector field $V = r/r^3$

- Q66. The surface integral of this vector over the surface of a cube of size a and centered at the origin
- (a) 0 (b) 2π (c) $2\pi a^3$ (d) 4π

- Q67. Which one of the following is not correct?
- (a) Value of the line integral of this vector around any closed curve is zero
 (b) This vector can be written as the gradient of some scalar function
 (c) The line integral of this vector from point P to point Q is independent of the path taken
 (d) This vector can represent the magnetic field of some current distribution

Data for Q. No. 68 to 69

Consider the motion of a particle in the potential $V(x)$ shown in the figure.



- Q68. Suppose the particle has a total energy $E = V_1$ in the figure. Then the speed of the particle is zero when it is at

- (a) point P (b) point Q (c) point S (d) point T

Q69. Which one of the following statements is NOT correct about the particle?

- (a) It experience no force when its position corresponds to the point Q on the curve
 (b) It experience no force when its position corresponds to the point R on the curve
 (c) Its speed is the largest when it is at S
 (d) It will be in a closed orbit between P and R if $E < V_1$

Data for Q. No. 70 to 71

A particle of mass m moving with speed v collides with a stationary particle of equal mass. After the collision, both the particles move. Let θ be the angle between the two velocity vectors

Q70. If the collision is elastic, then

- (a) θ is always less than 90° (b) θ is always equal to 90°
 (c) θ is always greater than 90° (d) θ cannot be deduced from the given data

Q71. If the collision is inelastic, then

- (a) θ is always less than 90° (b) θ is always equal to 90°
 (c) θ is always greater than 90° (d) θ could assume any value in the range 0° to 180°

Data for Q. No. 72 to 73

Consider two conducting plates of infinite extent, one plate at $z=0$ and the other at $z=L$, both parallel to the xy plane. The vector and scalar potential in the region between the plates is given by

$$A(r,t) = A_0 \hat{i} \cos(kx + \alpha) \cos(kct);$$

$$\phi(r,t) = 0$$

Q72. For this to represent a standing wave in the empty region between the plates:

- (a) $k = \frac{\pi}{L}$ and $\alpha = 0$ (b) $k = 2\pi/L$ and $\alpha = \pi/2$
 (c) $k = \pi/(2L)$ and $\alpha = \pi/2$ (d) $k = \pi/2L$ and $\alpha = 0$

Q73. The energy density at $z=0$ and $t=0$ is :

- (a) 0 (b) $\epsilon_0 c^2 k^2 A_0^2$
 (c) $(1/2)\mu_0 A_0^2 k^2$ (d) $(1/2)\mu_0 A_0^2 k^2 + (1/2)\epsilon_0 c^2 k^2 A_0^2$

Date for Q. No. 74 to 75

A particle is located in a three dimensional cubic well of width L with impenetrable walls.

Q74. The sum of the energies of the third and the fourth level is:

- (a) $10\pi^2 \hbar^2 / mL^2$ (b) $10\pi^2 \hbar^2 / 3mL^2$
 (c) $11\pi^2 \hbar^2 / 2mL^2$ (d) $15\pi^2 \hbar^2 / 2mL^2$

Q75. The degeneracy of the fourth level is given by

- (a) 1 (b) 2 (c) 3 (d) 4

Data for Q. No. 76 to 77

The normalized wave function ψ_1 and ψ_2 correspond to the ground state and the first excited state of a particle in a potential. You are given the information that the operator \hat{A} acts on the wave functions as $\hat{A}\psi_1 = \psi_2$ and $\hat{A}\psi_2 = \psi_1$

Q76. The expectation value of A for the state $\psi = (3\psi_1 + 4\psi_2) / 5$ is :
 (a) -0.32 (b) 0.0 (c) 0.75 (d) 0.96

Q77. Which of the following are eigenfunctions of \hat{A}^2 ?
 (a) ψ_1 and ψ_2 (b) ψ_2 and not ψ_1 (c) ψ_1 and not ψ_2 (d) neither ψ_1 nor ψ_2

Data for Q. No. 78 and Q.79

In the presence of an inhomogeneous weak magnetic field, spectral lines due to transitions between two sets of states were observed.

$$(1) {}^51s \rightarrow {}^5H_4 \text{ and } (2) {}^2D_{5/2} \rightarrow {}^2P_{3/2}$$

Q78. The types of Zeeman effect observed in (1) and (2) respectively are
 (a) normal, normal (b) anomalous, anomalous
 (c) anomalous, normal (d) normal, anomalous

Q79. The number of levels into which each of the above four terms split into respectively is:
 (a) 6, 4, 10, 8 (b) 4, 6, 10, 12 (c) 11, 9, 6, 4 (d) 9, 5, 12, 10

Data for Q.No. 80 to 82

A system consists of three spin-half particles, the z -components of whose spin $S_z(1)$, $S_z(2)$ and $S_z(3)$ can take value $+1/2$ and $-1/2$. The total spin of the system is $S_z = S_z(1) + S_z(2) + S_z(3)$

Q80. The total number of possible micro-states of this system is:
 (a) 3 (b) 6 (c) 7 (d) 8

Q81. The total number of micro-states with $S_z = 1/2$ is:
 (a) 3 (b) 5 (c) 6 (d) 7

Q82. Consider an ensemble of systems where each microstate has equal probability. The ensemble average of S_z is:
 (a) $-1/2$ (b) 0 (c) $1/2$ (d) $3/2$

Data for Q.No. 83 and 84

A gas of N particles is enclosed in volume V at a temperature T . The logarithm of the partition function is given

by $\ln Z = N \ln \left\{ (V - bN) (k_B T)^{3/2} \right\}$ where b is a constant with appropriate dimensions.

Q83. If P is the pressure of the gas, the equation of state is given by
 (a) $P(V - bN) = Nk_B T$ (b) $P(V - bN) = k_B T$

(c) $P(V - b) = Nk_B T$ (d) $P(V - bN) = k_B T$

Q84. The internal energy of the gas is given by

(a) $U = (1/2)k_B T$ (b) $U = Nk_B T$ (c) $U = (3/2)Nk_B T$ (d) $U = 2Nk_B T$

Data for Q. No. 85 to 86

A crystal belongs to a face centered cubic lattice with four atoms in the unit cell. The size of the crystal is 1 cm and its unit cell dimension is 1 nm . f is the scattering factor of the atom.

Q85. The number of atoms in the crystal is
 (a) 2×10^{21} (b) 4×10^{21} (c) 2×10^{23} (d) 4×10^{24}

Q86. The structure factors for $(0\ 1\ 0)$ and $(2\ 0\ 0)$ reflections respectively are
 (a) $2f$ and zero (b) zero and $4f$ (c) $2f$ and $2f$ (d) zero and zero

Data for Q. No. 87 to 88

An atomic bomb consisting of ${}^{235}\text{U}$ explodes and releases an energy of 10^{14} J . It is known that each ${}^{235}\text{U}$ which undergoes fission releases 3 neutrons and about 200 MeV of energy. Further, only 20% of the ${}^{235}\text{U}$

atoms in the bomb undergoes fission :

Q87. The total number of neutrons released is about


- (a) 4.7×10^{24} (b) 9.7×10^{24} (c) 1.9×10^{25} (d) 3.7×10^{25}

Q88. The mass of ^{235}U in the bomb is about

- (a) 1.5 kg (b) 3.0 kg (c) 6.1 kg (d) 12 kg

Data for Q. No. 89 to 90

The circuit below represents a non-inverting integrator



Q89. For high frequencies ($\omega \rightarrow \infty$) the input impedance is

- (a) 0 (b) R (c) $R/(1 + \omega RC)$ (d) ∞

Q90. For low frequencies ($\omega \rightarrow 0$) the input impedance is

(a) $V_0 = (1/RC) \int V_1 dt$

(b) The voltages at the inverting and non-inverting terminals of the op-amp are nearly

(c) The voltage at the non-inverting terminal of the op-amp and the current in the resistor attached to it are $\pi/2$ out of phase

(d) The current in the two resistors are in phase

ANSWER KEY

- | | | | | |
|---------|---------|---------|---------|---------|
| 1. (a) | 2. (d) | 3. (a) | 4. (d) | 5. (a) |
| 6. (c) | 7. (c) | 8. (b) | 9. (d) | 10. (b) |
| 11. (a) | 12. (b) | 13. (c) | 14. (a) | 15. (c) |
| 16. (b) | 17. (c) | 18. (a) | 19. (b) | 20. (b) |
| 21. (c) | 22. (b) | 23. (d) | 24. (a) | 25. (a) |
| 26. (b) | 27. (b) | 28. (d) | 29. (c) | 30. (a) |
| 31. (a) | 32. (b) | 33. (b) | 34. (b) | 35. (c) |
| 36. (c) | 37. (d) | 38. (c) | 39. (c) | 40. (c) |
| 41. (a) | 42. (c) | 43. (b) | 44. (b) | 45. (c) |
| 46. (a) | 47. (b) | 48. (c) | 49. (b) | 50. (c) |
| 51. (c) | 52. (a) | 53. (c) | 54. (a) | 55. (d) |
| 56. (a) | 57. (c) | 58. (a) | 59. (b) | 60. (b) |
| 61. (c) | 62. (d) | 63. (a) | 64. (c) | 65. (d) |
| 66. (d) | 67. (d) | 68. (a) | 69. (c) | 70. (b) |
| 71. (a) | 72. (b) | 73. (a) | 74. (a) | 75. (c) |
| 76. (c) | 77. (a) | 78. (b) | 79. (c) | 80. (d) |
| 81. (a) | 82. (b) | 83. (a) | 84. (c) | 85. (b) |
| 86. (b) | 87. (b) | 88. (c) | 89. (b) | 90. (a) |